

Quasi-stationary distributions for strongly Feller processes and application to hypoelliptic Hamiltonian systems

Boris NECTOUX, LMPB - Université Clermont Auvergne

In this talk, I will give a general framework [1] ensuring existence and uniqueness of quasi-stationary distributions for strongly Feller processes $(X_t, t \geq 0)$ on a set \mathcal{D} in the space of measures ν such that $\nu(W^{1/p}) < +\infty$, where W is a Lyapunov functional for the non-killed process $(X_t, t \geq 0)$ and $p > 1$. Exponential convergence (in this set of measures) of the law of the process (conditioned not to leave \mathcal{D}) towards the quasi-stationary distribution is also derived. These results are then applied to a wide range of hypoelliptic Hamiltonian systems $(X_t = (x_t, v_t), t \geq 0)$ in R^{2d} solution to

$$\begin{cases} dx_t = v_t dt, \\ dv_t = -\nabla V(x_t) dt - \gamma(x_t, v_t) v_t dt + \Sigma(x_t, v_t) dB_t, \end{cases} \quad (1)$$

when $\mathcal{D} = \mathcal{O} \times R^d$, $\mathcal{O} \subset R^d$. Such domains are indeed those of interest to justify the use of a kinetic Monte Carlo processes (also called Markov jump processes) to model the state-to-state dynamics of a molecular system whose evolution satisfies (1). The approach also applies to singular potentials V as the Lennard-Jones potential and the Coulomb potential [2].

-[1] *Quasi-stationary distribution for strongly Feller Markov processes by Lyapunov functions and applications to hypoelliptic Hamiltonian systems*. A. Guillin, B. Nectoux, L. Wu. 2020. Submitted.

-[2] *Quasi-stationary distribution for Hamiltonian dynamics with singular potentials*. A. Guillin, B. Nectoux, L. Wu. 2021. In preparation.